

The rings of Chariklo under close encounters with the giant planets

R.A.N. Araujo¹
ran.araujo@gmail.com

R. Sfair¹
rsfair@feg.unesp.br
and

O.C. Winter¹
ocwinter@gmail.com

ABSTRACT

The Centaur population is composed by minor bodies wandering between the giant planets and that frequently perform close gravitational encounters with these planets, which leads to a chaotic orbital evolution. Recently, the discovery of two well-defined narrow rings was announced around the Centaur 10199 Chariklo. The rings are assumed to be in the equatorial plane of Chariklo and to have circular orbits. The existence a well-defined system of rings around a body in such perturbed orbital region poses an interesting new problem. Are the rings of Chariklo stable when perturbed by close gravitational encounters with the giant planets? Our approach to address this question consisted of forward and backward numerical simulations of 729 clones of Chariklo, with similar initial orbits, for a period of 100 Myrs. We found, on average, that each clone suffers along its lifetime more than 150 close encounters with the giant planets within one Hill radius of the planet in question. We identified some extreme close encounters able to significantly disrupt or to disturb the rings of Chariklo. About 3 % of the clones lose the rings and about 4 % of the clones have the ring significantly disturbed. Therefore, our results show that in most of the cases (more than 90 %) the close encounters with the giant planets do not affect the stability of the rings in Chariklo-like systems. Thus, if there is an efficient mechanism that creates the rings, then these structures may be common among these kinds of Centaurs.

Subject headings: minor planets: individual (10199 Chariklo), planets and satellites: rings, planets and satellites: dynamical evolution and stability

1. Introduction

Among the orbits of the giant planets there is a population of small objects called Centaurs.

There is not a consensus on the definition of the Centaur population. According to the Minor

Planet Center - MPC/IAU, Centaurs are celestial bodies with perihelion beyond the orbit of Jupiter and with semi-major axes smaller than the semi-major axis of Neptune (MPC-UMPweb). Similarly, the JPL/NASA defines the Centaurs population as the objects with semi-major axes between 5.5 au and 30.1 au (JPLweb). Duffard et al. (2014) classify Centaurs as celestial bodies with orbits mostly in the region between Jupiter and

¹Univ - Universidade Estadual Paulista, Grupo de Dinâmica Orbital e Planetologia, CEP 12516-410, Guaratingueta, SP, Brazil.

TABLE 1
ORBITAL AND PHYSICAL PARAMETERS OF CHARIKLO AND ITS RINGS

Chariklo		Rings	
$a^{(1)}$	15.74 <i>au</i>		
$e^{(1)}$	0.171	Orbital radii (km) ⁽²⁾	390.6 ± 3.3
$i^{(1)}$	23.4°	Width (km) ⁽²⁾	7.17 ± 0.14
Equivalent radius (km) ⁽²⁾	124	Radial separation (km) ⁽²⁾	14.2 ± 0.2
Mass (kg) ⁽³⁾	7.986×10^{18}	Gap between rings (km) ⁽²⁾	8.7 ± 0.4
			404.8 ± 3.3
			$3.4^{+1.1}_{-1.4}$

⁽¹⁾Orbital elements obtained from JPL's Horizons system for the epoch MJD 56541. According to JPL the uncertainties in a , e and i are of the order 10^{-5} , 10^{-6} and 10^{-5} , respectively.
⁽²⁾ Braga-Ribas et al. (2014).
⁽³⁾Calculated considering a density of 1 g/cm^3 , the equivalent radius of Chariklo and a spherical body.

Neptune and that typically cross the orbits of the giant planets. A similar definition is also found in Horner et al. (2004b).

2060 Chiron was the first observed body of this population (Kowal, Liller & Marsden 1979). Since then, the number of known Centaurs has grown. Currently, more than 400 objects are cataloged (MPCweb), and from the flux of short period comets Horner et al. (2004a) estimated the total number of objects with diameter $> 1 \text{ km}$ to be approximately 44000.

The Centaur 10199 Chariklo was discovered in 1997 by the Spacewatch program (Spacewatchweb). In 2013, a stellar occultation revealed the existence of symmetric features encircling Chariklo, the second largest known Centaur. Braga-Ribas et al. (2014), showed that these structures are a system of two narrow and well-defined rings. These discovery made the debut of minor bodies within the select group of ringed objects.

In that work, the authors estimated that the rings have orbital radii of approximately 391 *km* and 405 *km*, and width of 7 *km* and 3 *km*, respectively. They are assumed to be in the equatorial plane of Chariklo, with circular orbits. Besides an equivalent radius of 124 *km* derived from the same stellar occultation, few information about the physical properties of Chariklo is available. From the orbital position of the rings, they also estimated the density of the central body to be 1 g/cm^3 . Table 1 summarizes some of the physical and orbital parameters of Chariklo and its rings.

A detailed study of the orbital evolution of the Centaurs was presented by Horner et al. (2004a), where they analyzed the orbital evolution of 32

cataloged objects through numerical simulations of an ensemble of particles under the influence of the Jovian planets. They followed the particles both forward and backward in time and registered the dynamical evolution and fate of the particles. The orbital radius shows that Chariklo orbits between Saturn and Uranus, corresponding to a typical *U* class object, i.e., those whose evolution is controlled by Uranus (Horner et al. 2003). For Chariklo, they found the half-life to be 10.3 (9.68) Myr.

The Centaur objects are transient. Therefore, a source is required to keep a steady state population. The idea of bodies coming from regions of the Solar System beyond Neptune and populating the region between the planets is well-accepted.

Levison & Duncan (1997) estimated, through numerical integrations, a number of $\approx 1.2 \times 10^7$ of comets transiting between the inner and outer Solar System originating from the Kuiper Belt. In fact, Horner et al. (2004a), estimated a flux of one body coming from the Kuiper belt and getting into the Centaurs population every 125 yr. Sisto & Brunini (2007), present the Scattered Disk Objects - SDO (bodies with distance to the perihelion $q < 30 \text{ au}$ and semi-major axis $a > 50 \text{ au}$) as the most probable source of the Centaurs. Emel'Yanenko et al. (2007), analyzed the role of the Oort cloud in determining the flux of cometary bodies through the planetary system. They concluded that a substantial fraction of all known cometary bodies may have a source in the Oort cloud, including the Centaur population, which they defined as the population small bodies with perihelion $5 < q < 28 \text{ au}$ and $a <$




Fig. 1.— Histograms of the fraction of Chariklo clones lost within 100 *Myrs*, as a function of time. a) Backward integration. b) Forward integration. Throughout the numerical integration the clones could be lost by ejection or collisions with one of the giant planets or with the Sun. Considering an ejection distance of 100 *au* and with collisions defined by the physical radius of the planets and of the Sun.

1000 *au*. Following the same definition for Centaurs, Emel'Yanenko et al. (2013) present that in fact more than 90% of all Centaurs with $a > 60$ *au*, and $\approx 50\%$ with $a < 60$ *au*, come from the Oort cloud. In Brassier et al. (2012), the Oort cloud is also pointed out as the source of Centaurs, especially those with high-inclination. In that work the Centaur population is defined as small bodies with perihelion between 15 and 30 *au* and semi-major axis shorter than 100 *au*. They showed that these objects probably were originated from the Oort cloud rather than the Kuiper belt or the scattered disc.

Along its orbital evolution a Centaur is strongly perturbed by the giant planets. Horner et al. (2004b), illustrate in detail the effects of those perturbations on the orbit of five selected Centaurs. The close encounters with the giant planets are quite frequent. As consequence, the Centaurs present a characteristic chaotic orbital evolution.

The existence of a small body owning a well-defined system of circular rings within such perturbed population poses an interesting new problem. This scenario has motivated the development of the present work. We investigated the stability

of the rings of Chariklo when perturbed by close encounters with the giant planets. We analyzed how effective are the close encounters in disturb or disrupt the rings of Chariklo. Furthermore, the development of this study may allow us to quantitatively evaluate how propitious the region of the Centaurs is for such small bodies owning their own systems of rings. A brief qualitative discussion on this subject is presented by Ortiz et al. (2015), where it is proposed that the Centaur Chiron may also have rings. The possible existence of two small bodies belonging to the same population and owning a system of rings is quite interesting, and indicate that such systems can be more frequent than expected.

Since our goal is to analyze the stability of the rings of Chariklo while perturbed by close encounter with the giant planets, here we classify it as a Centaur while its orbit is mainly in the region between Jupiter and Neptune (as in Duffard et al. (2014)), with maximum semi-major axis value of $a \leq 50$ *au*.

The structure of this paper is as follows. In Section 2, we present the initial conditions and the numerical method adopted in order to iden-

tify the close encounters of Chariklo with the giant planets. In Section 3, we present the selection of the extreme close encounters, i.e., those encounters that could be capable of disrupting the rings of Chariklo. In Section 4, we describe how the rings were simulated along a close encounter. In Secs. 5, 6 and 7 we present the results, and in Section 8, are our final comments and the major conclusions of the work.

2. Close encounters with the giant planets

The first step of the work consisted on selecting a representative sample of close encounters performed by Chariklo with the giant planets.

We considered a system composed by the Sun, the giant planets of the Solar System (Jupiter, Saturn, Uranus and Neptune), and a sample of clones, i.e., objects with the same mass and radius of Chariklo, but with small deviations on their orbits.

The clones were created following the procedure presented in Horner et al. (2004a), where 729 clones were created from the original orbit assuming a variation of semi-major axis of $0.005 au$, a variation of eccentricity of 0.005 and a variation of inclination of 0.01° .

The orbital elements of Chariklo and of the planets was obtained through JPL’s Horizons system for the epoch MJD 56541. For the orbit of Chariklo at this epoch we have $a = 15.74 au$, $e = 0.171$ and $i = 23.4^\circ$.

Considering these orbital elements, and taking the amplitude of variation as in Horner et al. (2004a) in such way that we have 729 clones, we created the clones of Chariklo orbiting the Sun as follows: $15.720 \leq a \leq 15.760 au$, taken every $0.005 au$; $0.151 \leq e \leq 0.191$, taken every 0.005 and, $23.36^\circ \leq i \leq 23,44^\circ$, taken every 0.01° . The choice of these values resulted in nine values of semi-major axes, nine values of eccentricities and nine values of inclination. The combination of these values resulted in 729 clones, each one with different values of a , e and i .

Considering that Chariklo has an equivalent radius of $124 km$ and a density of $1 g/cm^3$ (Braga-Ribas et al. 2014), we estimated its mass as $M_C = 7.986 \times 10^{18} kg$.

We performed backward and forward numerical integrations of the system composed by the Sun,

by the giant planets and by the clones, for a time span of $100 Myrs$, using the adaptive time-step hybrid symplectic/Bulirsch-Stoer algorithm from MERCURY (Chambers 1999).

Throughout the numerical integrations the clones did not interact with each other, but they could collide with the planets or escape from the system. The collisions were defined by the relative distance between the clones and the planets. If the clone-planet distance is smaller than the radius of the planet in question, then we have a collision. The physical radius of the planet is determined by MERCURY assuming a spherical planet with uniform density. We consider ejections as being the ejections from the Centaur population defined by the relative distance to the Sun of $100 au$. This value was adopted taking into account that if a clone reach the distance of $100 au$ and it is still in a elliptical orbit then necessarily the semi-major axis of the clone has to be greater than $50 au$, i.e, the clone is no longer classified as a Centaur, according to our definition.

As a result of the integrations, we see in the histograms in Figure 1 that more than 50% of our sample was lost (ejections or collisions) in $10 Myrs$, for both, backward and forward integrations. These results show that the evolution of our sample is in agreement with the predicted evolution of the Centaurs, which has an estimated mean lifetime of about $10 Myrs$ (Tiscareno & Malhotra 2003). We also note that there is a kind of symmetry on those results, which indicates that Chariklo is currently in the middle of its median dynamical lifetime as a Centaur.

At the end of the forward integrations, we found that $\approx 94\%$ of the 729 clones were lost in the time span of $100 Myrs$, being 683 clones lost by ejections and 4 clones lost by collisions (three with Saturn and one with Jupiter). For the backward integration, we found that $\approx 99\%$ of the clones were lost in $100 Myrs$, being 719 ejections and 4 collisions (three with Jupiter and one with Saturn).

Once we have characterized the evolution of the sample of clones as a whole, we then selected all close encounters of the clones within 1 Hill radius with each giant planet performed within $10 Myrs$ (mean lifetime of the Centaurs). For this time span, there were registered 60159 close encounters for the forward integration, and 65293 encounters

for the backward integration. From Table 2 we see that in this case Uranus dominates, followed by Saturn, Jupiter and Neptune, for both, backward and forward integrations. This result is in agreement with works on the dynamics of Centaurs stating that the dynamics of bodies with similar orbits to the orbit of Chariklo should be guided by Uranus, as discussed in Horner et al. (2004a).

However, we are interested in analyzing how the close encounters of Chariklo with the giant planets might affect its rings. Therefore, we selected among all these registered close encounters those that are expected to perturb or disrupt the rings. Following are the details of this analysis and the results obtained.

3. Extreme close encounters

In order to select the extreme close encounters, i.e., encounters with the giant planets that are expected to significantly affect the rings of Chariklo, we have calculated the Tidal Disruption Radius (r_{td}).

According to Philpott et al. (2010), the distance of the close encounter at which the tidal disruption of a binary may occur, is given by:

$$r_{td} \approx a_B \left(\frac{3M_{Pl}}{M_1 + M_2} \right)^{1/3} \quad (1)$$

where a_B is the semi-major axis of the binary, M_{Pl} is the mass of the planet and M_1 and M_2 are the masses of the components of the binary.

For a particle orbiting Chariklo with $a_B = 410$ km (approximated outer limit of the ring), we calculated the r_{td} for encounters performed with each one of the giant planets. These values are presented in Table 2.

It is important to point out that this is an approximated value since it does not take into account the relative velocity of the bodies at the moment of the encounter. Araujo et al. (2008), has shown that not just the distance of the encounter, but also the relative encounter velocity determine how significantly a body will be disturbed by a close encounter. Such effect were also discussed by Araujo & Winter (2012), when they compared their numerical analysis on the disruption of NEAs binaries due to close encounters with the Earth, with the analytical prediction given by

Eq. 1, showing the dependence of the results on the relative velocity of the encounters.

Nevertheless, for our purposes the approach given by Eq. 1 is adequate. Knowing that this value is an approximation, we then selected among all the registered close encounters those that had the minimum distance within $10 r_{td}$. For the forward integration, we see that most of them (about 3/4) occurred with Jupiter and Saturn (Table 2). Very few encounters occurred within $1r_{td}$ (none with Uranus or Neptune). For the backward integration, we see that the extreme encounters with Jupiter and Saturn still prevail, but here we note the occurrence of a few encounters occurred within $1r_{td}$ with Uranus and Neptune.

We explored the effects of each one of these extreme encounters ($\leq 10r_{td}$) on the particles of Chariklo's rings, as follows.

4. Simulating the rings

At the second step of the work, we numerically simulated the extreme close encounters including massless particles around Chariklo.

According to Section 2, a close encounter is registered when a clone of Chariklo crosses the limit of 1 Hill radius of any of the giant planets. At this crossing moment, we recorded the position and the velocity of these bodies, relative to the Sun. These values are the initial conditions for the simulations of the extreme close encounters, i.e., encounters with minimum distance within $10r_{td}$. Thus, at this step, the numerical simulations always involve the Sun, the bodies performing the close encounter (planet and Chariklo), and a sample of particles orbiting Chariklo.

We considered particles with circular equatorial orbits, radially distributed from 200 km to 1000 km, taken every 20 km. For each radial distance there were considered 100 particles in a random angular distribution. Such combination of values resulted in a total of 4100 particles orbiting Chariklo.

The pole orientation of Chariklo was considered perpendicular to the orbital plane. This is a reasonable approach since the rings are assumed to be in the equatorial plane of Chariklo, and we are interested in the maximized radial perturbations of the rings.

The encounters were simulated for a time span

TABLE 2

REGISTERED CLOSE ENCOUNTERS OF THE CLONES WITH EACH ONE OF THE GIANT PLANETS WITHIN 1 HILL RADIUS AND WITHIN 1 AND 10 RUPTURE RADIUS (r_{td}), IN THE TIME SPAN OF 10 *Myrs*, FOR BOTH, FORWARD AND BACKWARD INTEGRATIONS.

Planet	Hill Radius ⁽¹⁾ (Planetary radii)	r_{td} ⁽¹⁾ (Planetary radii)	Forward Encounters			Backward Encounters		
			1 Hill Radius ⁽²⁾	$r_{td} \leq 1$	$1 < r_{td} \leq 10$	1 Hill Radius ⁽³⁾	$r_{td} \leq 1$	$1 < r_{td} \leq 10$
Jupiter	≈ 740	≈ 5	16.6%	5	36	18.3%	5	47
Saturn	≈ 1100	≈ 4	26.0%	1	34	24.2%	0	25
Uranus	≈ 2700	≈ 5	48.0%	0	18	46.9%	2	13
Neptune	≈ 4600	≈ 5	9.4%	0	2	10.6%	1	5

⁽¹⁾ The Hill radius and the rupture radius in terms of the radius of the planet in question.

⁽²⁾Percentage relative to 60159 encounters.

⁽³⁾Percentage relative to 65293 encounters.

TABLE 3


REGISTERED CATASTROPHIC AND DISTURBING ENCOUNTERS OF THE RINGS OF CHARIKLO DUE TO EXTREME CLOSE ENCOUNTERS WITH EACH ONE OF THE GIANT PLANETS, IN THE TIME SPAN OF 10 MYRS.

	Forward Encounters			Backward Encounters		
	Catastrophic ⁽¹⁾	Disturbing ⁽²⁾	Survival Time ⁽³⁾ [Max:Min](years)	Catastrophic ⁽¹⁾	Disturbing ⁽²⁾	Survival Time ⁽³⁾ [Max:Min](years)
Jupiter	6	9	[16,125:221,739]	6	16	[-49,245:-546,550]
Saturn	4	7	[56,650:623,559]	0	7	-
Uranus	0	0	-	3	1	[-1,371,579:-4,401,849]
Neptune	0	0	-	1	2	-1,499,269


⁽¹⁾ Rings are completely removed.

⁽²⁾ External particles removed but the rings survive.

⁽³⁾ Minimum and maximum survival time registered among the clones after they suffered a catastrophic encounter. excluding the immediate ejection cases.



(a)



(b)

Fig. 2.— Example of a disturbing close encounter caused on the ring by a close encounter with Jupiter. a) Maximum final change in eccentricity. b) Maximum final change in semi-major axis (km). The plots show the maximum final of those elements among the hundred particles that share the same initial radial position. The vertical lines indicate the boundary of the rings’ region. The encounter were performed with a minimum distance of 6.6 planet radius, with a relative velocity $V_\infty = 8.00$ km/s. The encounter resulted in the critical radius of $R_C = 460$ km, meaning that no particle in the region of the rings were removed.

of 1 year using the adaptive time-step Gauss-Radau numerical integrator, keeping the accuracy 10^{-12} (Everhart 1985). Throughout the integration the particles could collide with Chariklo or be ejected. The collisions were defined by the equivalent radius of Chariklo (124 km). The ejections were defined by the energy of the Two-body Problem Chariklo-particle.

According to Table 2, there were simulated 96 close encounters for the forward case, and 98 for the backward case. The results of these simulations and their implications are presented in Table 3, and they are discussed in the following sections.

5. Catastrophic Encounters

We classify as catastrophic those close encounters that lead to the complete removal of the particles in the region of the rings of Chariklo. Knowing that the particles of the rings are distributed in the range of ≈ 390 km to ≈ 405 km, we defined that there was a catastrophic encounter if at the end of our simulation particles distributed beyond 380 km were lost by ejection or collision as defined in Sec. 4.

The results presented in Table 3 show that in about 10% of the simulations the rings were removed from Chariklo due to close gravitational encounters with the giant planets, for both, backward and forward integrations.

For the forward integration, we found that only extreme encounters with Jupiter and Saturn were able to fully remove the rings. For the backward integration there were a few cases where Uranus and Neptune were able to do that. Our data suggest that Uranus and Neptune might have influenced the existence of the rings in the past, but from now on Jupiter and Saturn would play this role. However, more simulations are required to investigate whether this is a real difference in the forward/backward evolution or the result of statistical artifact due to the small number of extreme close encounters.

In the cases of forward integration in time, after the removal of the rings, Chariklo remained as a Centaur for much less than a million years, i.e., the rings were destroyed in the very last stage of Chariklo’s orbital evolution among the giant planets.

In the cases of backward integration in time, the removal of the rings occurred much before Chariklo complete its first million years as a Centaur, i.e., in its first stage of orbital evolution among the giant planets.

6. Disturbing Encounters

There were cases in which the particles of the rings were not removed, but their orbits were significantly changed due to a significant perturbation caused by the encounter with a giant planet.

In Figure 2, we present an example of the effects of such kind of encounter with Jupiter. The plots show the maximum final variation in semi-major axis and eccentricity among the hundred particles that share the same initial radial position. As expected, the values increase with the radial distance. In the region of the rings (indicated by the vertical lines) the semi-major axis changes by more than 30 km and the eccentricity grows up to more than 0.5.

The results of the encounters that produced at least some noticeable variations ($\Delta e \geq 0.01$) on the orbits of the rings are presented in Table 4, for backward and forward integrations, excluding the encounters followed by ejection.

For the forward integration, we observed only five encounters with Jupiter and two with Saturn that resulted in eccentricities larger than 0.1 for the ring particles. None of the encounters with Uranus and Neptune showed significant orbital variation on the particles of the rings. Similarly, if we look backward, we see fewer close encounters that could have increased the eccentricity of the particles of the rings for values larger than 0.1 (5 with Jupiter, 1 with Saturn and 1 with Uranus).

Since Chariklo, as observed now, has well-defined narrow circular rings (Braga-Ribas et al. 2014), it might mean that it did not suffer any of those disturbing encounters or if it happened it was so long ago so that the rings had time to evolve damping the eccentricity and reshaping them. Both possibilities are compatible with our results, and as shown in Table 3, the probability of catastrophic or disturbing encounters is very low, and probably occurred several million years ago.

7. The ejection cases

The close encounters of Chariklo with any of the giant planets followed by ejections are especially interesting. If we look forward, those would be the encounters responsible for removing Chariklo from the population of Centaurs. On the other hand, looking backward, those events would be the ones that brought Chariklo into the region that we defined as the Centaurs population, according to our definition (see Sec. 1). We then looked for the last close encounter (within 1 Hill radius) that every clone suffered before being ejected.

For the forward integration, we found that approximately 36% of the clones suffered an encounter with Jupiter before the ejection, 45% with Saturn, 10% with Neptune, and 9% with Uranus.

For the backward integration, we found that approximately 27% of the encounters were with Jupiter, 42% with Saturn, 20% with Neptune, and 11% with Uranus.

As expected, the same pattern is obtained when we restrict our analysis to the total of extreme close encounters, i.e., encounters performed within $1 \leq r_{td} \leq 10$ (Table 2 and Table 3). We found for the forward integrations that 4 clones of Chariklo were ejected after an extreme close encounter with Jupiter, 3 with Saturn, and none with Uranus or Neptune. For the backward integrations, we registered 11 clones ejected after a close encounter with Jupiter, 5 with Saturn, 2 with Uranus, and 1 with Neptune.

Among these encounters, we found that only in 6 cases Chariklo was ejected just by the close encounter that disrupted the rings. For the forward integrations, such kind of event occurred 2 times after an encounter with Jupiter and one time after an encounter with Saturn. For the backward integrations, we registered 2 cases with Jupiter and one with Uranus.

Thus, our results indicate that for both backward and forward integrations, Jupiter and Saturn are the mainly responsible for inserting and removing Chariklo from the Centaurs region.

However, it is important to point out that since our ejection criteria requires that the object goes beyond 100 *au*, it is more likely to have ejections caused by stronger close encounters as those produced by Jupiter and Saturn. Another con-

TABLE 4

LIST OF SIGNIFICANT DISTURBING ENCOUNTERS ($\Delta e \geq 0.01$) WITH THE GIANT PLANETS FOR BOTH, FORWARD AND BACKWARD INTEGRATION WITHIN THE TIME SPAN OF 10 MYRS. THE IMMEDIATE EJECTION CASES WERE EXCLUDED.

Jupiter - Forward				
Minimum encounter distance (Planet radius)*	V_∞ (km/s)	Maximum ring Δa (km)	Maximum ring Δe	Survival Time** (years)
6.6	4.00	35.28	0.54	451,203
6.9	4.67	7.61	0.28	509,333
8.5	5.81	34.40	0.45	43,868
8.8	3.76	28.20	0.32	6,891
9.2	5.03	44.33	0.57	1,140,000
15.4	7.00	0.04	0.02	55,034
16.0	7.78	0.02	0.01	59,483
Saturn - Forward				
6.8	4.87	24.77	0.34	22,115
7.4	3.12	30.13	0.53	337,731
8.2	4.42	1.23	0.07	87,594
9.3	5.29	0.68	0.06	253,520
10.4	3.63	0.14	0.031	573,463
Jupiter - Backward				
4.2	9.87	25.59	0.20	-217,796
8.3	4.17	17.34	0.40	-18,259
8.4	4.96	13.63	0.33	-23,770
8.4	10.32	65.85	0.51	-8,204
8.5	7.23	2.42	0.20	-280,371
11.1	4.93	0.065	0.03	-84,066
11.1	5.20	5.74	0.19	-153,800
11.9	6.62	2.33	0.10	-537,024
13.4	3.17	0.28	0.04	-309,943
13.5	7.40	0.06	0.02	-36,896
14.4	1.25	0.03	0.01	-98,210
15.4	5.72	0.02	0.01	-45,522
16.7	9.65	0.03	0.01	-1,634,011
Saturn - Backward				
7.9	2.68	13.79	0.300	-405,570
Uranus - Backward				
10.8	3.00	3.73	0.11	-2,504,843

* The minimum encounter distance given in terms of the radius of the planet in question.
** Survival time after a disturbing encounter.

sequence of our ejection criteria is that there are bodies that became TNOs without being considered ejected. For example, a Plutino-like object with $a = 39 \text{ au}$ and $e = 0.2$ would remain as a Centaur in our simulations.

Anyhow, regardless the way that Chariklo followed to get in or to go out from the Centaur region, our results allow us to conclude that it may have brought the rings when it was introduced into this region and that it can keep them while leaving it.

8. Final Comments

In the present work we explored the dynamics of the Centaur 10199 Chariklo and the stability of its rings when disturbed by frequent close encounters with the giant planets.

Through numerical integrations we analyzed the orbital evolution of Chariklo while a Centaur (orbits mainly in the region between Jupiter and Neptune and with $a \leq 50 \text{ au}$).

To do so we considered a sample of 729 clones of Chariklo with small deviations on semi-major axis, eccentricity and inclination.

Throughout the lifetime of Chariklo in the population of the Centaurs, we found that the close encounters within 1 Hill radius with Uranus are more frequent than the encounters with the other planets. Nevertheless, if we look to the most significant close encounters, i.e, those able to disturb or disrupt the rings of Chariklo, we found that Jupiter and Saturn dominate.

For the forward integrations the most significant encounters happened in the last stage of Chariklo among the giant planets, and they were performed exclusively with Jupiter and Saturn. This indicates that Uranus and Neptune may not play an important role in the future dynamics of the rings of Chariklo, but these planets may have had more influence on the rings in the past. For the backward integrations we found few cases of catastrophic encounters of Chariklo with these planets, being three catastrophic encounters with Uranus and one with Neptune. Nevertheless, the difference between forward and backward integrations is subtle and probably arised from a statistical analysis based on a small number of extreme close encounters.

In total, for both backward and forward inte-

grations, we found that the number of close encounters really able to completely disrupt the rings of Chariklo is small ($\approx 3\%$ of the clones), and they most probably happen in the beginning or in the final stages of Chariklo's time as a Centaur.

Thus, although a typical Centaur such as Chariklo present a chaotic orbital evolution (Horner et al. 2004a), we found that if the body had those rings before becoming a Centaur or acquired them in the region among the giant planets, then these rings will probably survive throughout its Centaur life. Hence, the formation of the rings of Chariklo while a Centaur is not mandatory.

Therefore, our major conclusion is that Centaurs experience a propitious environment to the existence of rings. Furthermore, if there is an efficient mechanism that creates the rings, then these structures may be common among the bodies of this population.

One possible mechanism to create rings around Centaurs could be the outcome of extreme close encounters with the giant planets obtained here. This possibility will be analyzed in a future work.

9. Acknowledgments

This work was funded by CAPES, CNPq and FAPESP (proc. 2011/08171-3). This support is gratefully acknowledged. The authors are also grateful to the anonymous referee for suggestions that significantly improved the paper.

REFERENCES

- Araujo, R.A.N.; Winter, O.C.; Prado, A.F.B.A.; Vieira Martins, R. Sphere of Influence and Gravitational Capture Radius: A Dynamical approach. *MNRAS*, v. 391, Issue 2, p.675-684, 2008.
- Araujo, R. A. N.; Winter, O. C. Near-Earth asteroid binaries in close encounters with the Earth. *Astronomy & Astrophysics*, 566, id.A23, 10 pp., 2012.
- Braga-Ribas,F, et al. A ring system detected around the Centaur (10199) Chariklo. *Nature*, Vol. 508, Issue 7494, pp. 72-75, 2014.
- Brasser, R.; Schwamb, M. E.; Lykawka, P. S.; Gomes, R. S. An Oort cloud origin for the high-

- inclination, high-perihelion Centaurs. *MNRAS*, 420, Issue 4, pp. 3396-3402, 2012.
- Chambers, J.E. A hybrid symplectic integrator that permits close encounters between massive bodies. *MNRAS*, 304, pp. 793-799, 1999.
- Duffard, R.; Pinilla-Alonso, N.; Santos-Sanz, P.; Vilenius, E.; Ortiz, J. L.; Mueller, T.; Fornasier, S.; Lellouch, E.; Mommert, M.; Pal, A.; Kiss, C.; Mueller, M.; Stansberry, J.; Delsanti, A.; Peixinho, N.; Trilling, D. TNOs are Cool”: A survey of the trans-Neptunian region. XI. A Herschel-PACS view of 16 Centaurs. *A&A*, Volume 564, id.A92, 17 pp, 2014.
- Emel’Yanenko, V. V.; Asher, D. J.; Bailey, M. E. The fundamental role of the Oort cloud in determining the flux of comets through the planetary system. *MNRAS*, 381, Issue 2, pp. 779-789, 2007.
- Emel’Yanenko, V. V.; Asher, D. J.; Bailey, M. E. A Model for the Common Origin of Jupiter Family and Halley Type Comets. *Earth, Moon, and Planets*, 110, Issue 1-2, pp. 105-130, 2013.
- Everhart, E. An efficient integrator that uses Gauss-Radau spacings. In *Dynamics of comets: Their origin and evolution*, Eds. A. Carusi Carusi and G. B. Valsecchi, D.Reidel Publishing Company (Holanda), pp. 185-202, 1985.
- Horner, J.; Evans, N. W.; Bailey, M. E.; Asher, D. J. The populations of comet-like bodies in the Solar system. *MNRAS*, 343, Issue 4, pp. 1057-1066, 2003.
- Horner, J.; Evans, N. W.; Bailey, M. E. Simulations of the population of Centaurs - I. The bulk statistics. *MNRAS*, 354, Issue 3, pp. 798-810, 2004a.
- Horner, J.; Evans, N. W.; Bailey, M. E. Simulations of the population of Centaurs - II. Individual objects. *MNRAS*, Vol. 355, Issue 2, pp. 321-329, 2004b.
- JPLweb - Jet Propulsion Laboratory. Available in http://ssd.jpl.nasa.gov/sbdb_help.cgi?class=CEN. Accessed in: 4th February, 2016.
- Kowal, C. T.; Liller, W.; Marsden, B. G. The discovery and orbit of 2060 Chiron. In: *Dynamics of the solar system; Proceedings of the Symposium, Japan, Dordrecht*, D. Reidel Publishing Co., 1979, p. 245-250.
- Levison, H. F.; Duncan, M. J. From the Kuiper Belt to Jupiter-Family Comets: The Spatial Distribution of Ecliptic Comets. *Icarus*, Vol. 127, Issue 1, pp. 13-32, 1997.
- MPCweb - Minor Planet center. Available in <http://www.minorplanetcenter.net/iau/lists/Centaurs.html>. Accessed in: 18th November, 2015.
- MPC-UMPweb - Minor Planet center - Unusual Minor Planets. Available in <http://www.minorplanetcenter.org/iau/lists/Unusual.html>. Accessed in: 4th February, 2016.
- Ortiz, J. L.; Duffard, R.; Pinilla-Alonso, N.; Alvarez-Candal, A.; Santos-Sanz, P.; Morales, N.; Fernandez-Valenzuela, E.; Licandro, J.; Campo Bagatin, A.; Thirouin, A. Possible ring material around Centaur (2060) Chiron. *A&A*, Vol. 576, id.A18, 12 pp.
- Philpott, C. M.; Hamilton, D. P.; Agnor, C. B. Three-body capture of irregular satellites: Application to Jupiter. *Icarus*, Vol. 208, Issue 2, pp. 824-836, 2010.
- Sisto, R.P.; Brunini, A. The origin and distribution of the Centaur population. *Icarus*, Vol. 190, Issue 1, pp. 224-235, 2007.
- Spacewatchweb - Spacewatch Program. Available in <http://spacewatch.lpl.arizona.edu/discovery.html>. Accessed in: 18th November, 2015.
- Tiscareno, M.S, Malhotra, R. The Dynamics of known Centaurs. *The Astronomical Journal*, 126, pp. 3122-3131, 2003.